

Abstract Submitted  
for the MAR09 Meeting of  
The American Physical Society

**Momentum density, Fermi surface and directional Compton profile in the Iron-based superconductor LaOFeAs** Y. J. WANG, HSIN LIN, B. BARBIELLINI, Northeastern U., P.E. MIJNARENDS, Northeastern U. and U. Tech., Delft, The Netherlands, S. KAPRZYK, Northeastern U. and Academy of Mining and Metallurgy AGH, Poland, W. AL-SAWAI, R.S. MARKIEWICZ, A. BANSIL, Northeastern U. — We have carried out first principles all-electron calculations of the (001) projected 2D electron momentum density (2D-EMD) and directional Compton profiles (CPs) along the [100], [001], and [110] directions in iron-based superconductor  $\text{LaO}_{1-x}\text{F}_x\text{FeAs}$  for various doping concentrations  $x$  within the framework of the local density approximation (LDA). We have identified Fermi surface features both in the 2D-EMD and in the CPs. Bonding effects related to the character of wave functions near the Fermi level are revealed by the autocorrelation function  $B(\mathbf{r})$  defined as the Fourier transform of the momentum density. Work supported in part by the US Department of Energy.

Yung Jui Wang  
Northeastern U.

Date submitted: 19 Nov 2008

Electronic form version 1.4