

Abstract Submitted
for the MAR09 Meeting of
The American Physical Society

Non-equilibrium tunneling spectroscopy in carbon nanotubes¹

NADYA MASON, YUNG-FU CHEN, TRAVIS DIRKS, University of Illinois at Urbana-Champaign, GASSEM AL-ZOUBI, NORMAN BIRGE, Michigan State University — We report measurements of the non-equilibrium electron energy distributions in carbon nanotubes. Carbon nanotubes can be considered model one-dimensional systems whose transport is strongly affected by electron interactions. Using tunneling spectroscopy via a superconducting probe, we have studied electron energy distribution functions, and hence inelastic electron scattering, in nanotubes that have bias voltages applied between their ends. We find that at low temperatures, electrons interact weakly in nanotubes of a few microns channel length, independent of end-to-end conductance values. Surprisingly, the energy relaxation rate can increase substantially when the temperature is raised to only 1.5 K. In general, tunneling spectroscopy with a superconducting probe may be a powerful new tool for characterizing electron behavior in carbon nanotubes.

¹Work supported by the DOE under DE-FG02-07ER46453 through the FS-MRL and the NSF under DMR-0405238 and 0705213.

Nadya Mason
University of Illinois

Date submitted: 20 Nov 2008

Electronic form version 1.4