

Abstract Submitted
for the MAR09 Meeting of
The American Physical Society

Electrical detection of dynamic nuclear polarization and nuclear magnetic resonance in ferromagnet-semiconductor heterostructures¹

MUN CHAN, J. ZHANG, University of Minnesota, Twin Cities, Q. HU, University of California, Santa Barbara, T. KONDO, E. GARLID, University of Minnesota, Twin Cities, C.J. PALMSTROM, University of California, Santa Barbara, P.A. CROWELL, University of Minnesota, Twin Cities — We report all-electrical measurements of dynamic nuclear polarization (DNP) and nuclear magnetic resonance (NMR) in ferromagnet-semiconductor heterostructures. Nuclei in GaAs are polarized by electron spins injected through an epitaxial Fe/GaAs Schottky tunnel barrier. In an oblique Hanle geometry, electron spin depolarization due to the hyperfine field is detected by measuring the change in spin-dependent electrochemical potential with a Fe contact. We also report electrical measurements of both magnetic and current driven NMR. A transverse magnetic field or the injection current is modulated at the NMR frequency to suppress DNP, resulting in changes of up to 80% in the spin-dependent voltage. We measured resonant frequencies of different isotopes in the GaAs channel. Harmonics of the fundamental transitions are observed, reflecting the existence of either quadrupolar coupling or dipole-dipole interactions.

¹This work was supported by the Office of Naval Research and NSF under DMR 08-04244.

M.K. Chan
University of Minnesota, Twin Cities

Date submitted: 21 Nov 2008

Electronic form version 1.4