

Abstract Submitted
for the MAR09 Meeting of
The American Physical Society

Spin correlations and magnetic excitation spectrum of electron-doped $\text{Nd}_{2-x}\text{Ce}_x\text{CuO}_{4\pm\delta}$ near the magnetic quantum critical point EUGENE MOTOYAMA, GUICHUAN YU, YUAN LI, Stanford Univ., DANIEL PETITGRAND, LLB, France, KLAUDIA HRADIL, Univ. Gottingen, Germany, RICHARD MOLE, Forschungsneutronenquelle Heinz Maier-Leibnitz, Germany, PATRICK MANG, INNA VISHIK, Stanford Univ., OWEN VAJK, Univ. of Missouri-Columbia, MARTIN GREVEN, Stanford University — One of the most intriguing issues in the field of high- T_c superconductivity is the electron-hole asymmetry: the hole- or electron-doping of the parent Mott insulators leads to superconductors with differing properties. For the comparatively less-studied electron-doped materials, the antiferromagnetic phase extends much further with doping and appears to overlap with the superconducting phase. Our previous inelastic neutron scattering measurements show that in the compound $\text{Nd}_{2-x}\text{Ce}_x\text{CuO}_{4\pm\delta}$, genuine long-range antiferromagnetism does not extend as far as previously thought, and may not coexist with bulk superconductivity; the system features a magnetic quantum critical point at $x\sim 0.13$, very close to the composition above which superconductivity is first observed [Motoyama *et al.*, Nature **445**, 186 (2007)]. Here we present new measurements for the instantaneous spin correlations and the magnetic excitation spectrum that aim to refine our understanding of the physics in this interesting doping regime.

Eugene Motoyama
Stanford Univ.

Date submitted: 20 Nov 2008

Electronic form version 1.4