## Abstract Submitted for the MAR09 Meeting of The American Physical Society

Dzyloshinskii-Moriya interactions in valence bond systems II<sup>1</sup> MAYRA TOVAR, KUMAR RAMAN, KIRILL SHTENGEL, UC Riverside — We investigate the effect of Dzyaloshinskii-Moriya interactions on the low temperature magnetic susceptibility for a system whose low energy physics is dominated by short-range valence bonds (singlets). Our general perturbative approach is applied to specific models expected to be in this class, including the Shastry-Sutherland model of the spin-dimer compound  $SrCu_2(BO_3)_2$  and the antiferromagnetic Heisenberg model of the recently discovered S=1/2 kagome compound  $ZnCu_3(OH)_6Cl_2$ . The central result is that a short-ranged valence bond phase, when perturbed with Dzyaloshinskii-Moriya interactions, will remain time-reversal symmetric in the absence of a magnetic field but the susceptibility will be nonzero in the zero temperature limit. Applied to  $ZnCu_3(OH)_6Cl_2$ , this model provides an avenue for reconciling experimental results, such as the lack of magnetic order and lack of any sign of a spin gap, with known theoretical facts about the kagome Heisenberg antiferromagnet.

 $^{1}$ NSF DMR-0748925

Mayra Tovar UC Riverside

Date submitted: 20 Nov 2008 Electronic form version 1.4