

Abstract Submitted
for the MAR09 Meeting of
The American Physical Society

Photoinduced EPR study of electron traps in TiO₂ crystals: Oxygen vacancies and Ti³⁺ ions SHAN YANG, ADAM BRANT, LARRY HALBURTON, Physics Department, West Virginia University — Electron paramagnetic resonance (EPR) provides a sensitive method to monitor native defects in wide-band-gap semiconductors. In-situ illumination with laser light at low temperature (photoinduced EPR) forms paramagnetic defects in fully oxidized bulk TiO₂ crystals. Illumination with 442 nm laser light at 30 K and below produces four electronlike centers and one holelike center. Three of the electronlike centers have $S = 1/2$ and are assigned, respectively, to a substitutional Ti³⁺ ion in the otherwise perfect lattice, a substitutional Ti³⁺ ion adjacent to a Si⁴⁺ ion, and a substitutional Ti³⁺ ion adjacent to an oxygen vacancy. The fourth electronlike center has $S = 1$ and is assigned to two Ti³⁺ ions adjacent to one oxygen vacancy. The holelike center has $S = 1/2$ and consists of a hole shared equally by two adjacent oxygen ions in the otherwise perfect lattice. Spin-Hamiltonian parameters, obtained from complete sets of angular dependence data, are presented for each of the centers. This work was supported by NSF Grant No. DMR-0804352.

Shan Yang
Physics Department, West Virginia University

Date submitted: 21 Nov 2008

Electronic form version 1.4