Abstract Submitted for the MAR09 Meeting of The American Physical Society

Relaxation of Terrace-width Distributions: Novel Analysis and Features¹ T.L. EINSTEIN, Univ. of Maryland, College Park (UM), AJMI BH. HAMOUDA, UM & Univ. Monastir, Tunisia, A. PIMPINELLI, UM & U.B.P.Clermont-2 & Science Attache, French Consulate, Houston — We describe a Fokker-Planck scheme to describe the relaxation of the terrace-width distribution (TWD) on a vicinal surface toward the generalized Wigner form describing equilibrium.² We performed KMC calculations on the standard minimal SOS model to show that the time constant gives physical information, in particular the energy barrier of the rate-determining process. For close-packed steps, this involves kinkantikink generation, breaking 3 rather than the expected 2 lateral bonds (the latter associated with equilibrium fluctuations). We discuss strengths and limitations of this FP procedure, higher moments of the distribution beyond the variance, and generalizations to other step orientations.

¹Work at UM supported by the MRSEC, NSF Grant DMR 05-20471 ²A. BH. Hamouda, A. Pimpinelli, & TLE, Surface Sci. 602 (2008) 3569

> Theodore Einstein University of Maryland, College Park

Date submitted: 21 Nov 2008

Electronic form version 1.4