

Abstract Submitted  
for the MAR09 Meeting of  
The American Physical Society

**Evidence for two gaps from specific heat in LiFeAs single crystals**

FENGYAN WEI, Texas Center for Superconductivity at University of Houston and Department of Physics, BING LV, Texas Center for Superconductivity at University of Houston and Department of Chemistry, FENG CHEN, YUYI XUE, Texas Center for Superconductivity at University of Houston and Department of Physics, ARNOLD M. GULOY, Texas Center for Superconductivity at University of Houston and Department of Chemistry, CHINGWU CHU, Texas Center for Superconductivity at University of Houston, Lawrence Berkeley National Laboratory, Hong Kong University of Science and Technology — The mechanism and the pairing state of the newly discovered FeAs based superconductors are still open questions. We report here the specific heat of several single crystals of LiFeAs from 1.8K to 160K. The  $C_p$  anomaly around the transition temperature  $T_c = 17$  K is clear, and a change of the slope  $d(C_p/T)/dT^2$  is observed around 4 K. This down turn in the  $C_p/T$  vs.  $T^2$  plot, however, is suppressed in both poly-crystal samples and under high fields. We attribute this feature to the second gap, and will compare the data with calculations.

Fengyan Wei  
Texas Center for Superconductivity  
and Physics Department, University of Houston

Date submitted: 10 Dec 2008

Electronic form version 1.4