

Abstract for an Invited Paper
for the MAR09 Meeting of
The American Physical Society

Dynamic nuclear polarization with single electron spins¹

JACOB TAYLOR, Massachusetts Institute of Technology

Hyperfine interactions limit electron spin coherence times in GaAs quantum dots. By separating a spin singlet state on a chip, we measure an ensemble averaged spin dephasing time T_2^* of 10 ns, limited by the contact hyperfine interaction with the GaAs host nuclei². We use electrical control of the exchange interaction to drive coherent spin rotations. Exchange driven spin rotations are used to implement a “singlet-triplet spin echo” pulse sequence, which leads to a spin coherence time, T_2 , exceeding 1 microsecond. We show that nuclear spins can be polarized by controlling two-electron spin states near the anti-crossing of the singlet (S) and triplet (T_+). An initialized S state is cyclically brought into resonance with the T_+ state, where hyperfine fields drive rapid rotations between S and T_+ , “flipping” an electron spin and “flopping” a nuclear spin³. The resulting Overhauser field approaches 80 mT, in agreement with a simple rate-equation model. A self-limiting pulse sequence is developed that allows the steady-state nuclear polarization to be set using a gate voltage.

¹In collaboration with J. Petta, M. Kolodrubetz, A. C. Johnson, A. Yacoby, M. D. Lukin, C. M. Marcus, M. P. Hanson, A. C. Gossard. Financially supported by the Sloan Foundation, the David and Lucille Packard Foundation, and NSF DMR-0819860.

²J. R. Petta *et al.*, *Science* **309**, 2180 (2005).

³J. R. Petta, J. M. Taylor *et al.*, *Phys. Rev. Lett.* **100**, 067601 (2008).