

Abstract Submitted
for the MAR09 Meeting of
The American Physical Society

Tunneling Spectroscopy of Ultrathin Insulating Films: Cu₂N on Cu(100)¹ CHARLES RUGGIERO, TAEYOUNG CHOI, JAY GUPTA, The Ohio State University — Insulating films of only a few atomic layers offer insight into the evolution of electronic structure at the nanoscale. We report scanning tunneling microscopy (STM) studies of one monolayer Cu₂N films grown on Cu(100). Our tunneling spectra indicate that Cu₂N acts as an insulator, with a band gap that exceeds 4 eV [1]. We study changes in this electronic structure with size, ranging from few-atom islands to complete films. We find that the conduction band edge first emerges in few-atom islands, and shifts toward lower energy with increasing island size. Images of the local density of states show standing wave patterns consistent with the confinement of electrons to these 2D islands. Measurements of the tunneling barrier height and image potential states indicate that the Cu₂N work function is ~0.9 eV larger than bare Cu. This suggests a significant surface dipole, consistent with charge transfer predicted by theory. <http://www.physics.ohio-state.edu/~jgupta>
[1] C.D. Ruggiero, T. Choi, J.A. Gupta, Appl. Phys. Lett. **91**, 253106 (2007).

¹NSF CAREER (DMR-0645451).

Charles Ruggiero
The Ohio State University

Date submitted: 26 Nov 2008

Electronic form version 1.4