

Abstract Submitted
for the MAR09 Meeting of
The American Physical Society

Rotating Bose-Einstein condensates at the phase transition point

MAHIR HUSSEIN, Max Planck Institut für Physik Komplexer Systeme, Dresden, Germany, PIET VAN ISACKER, Grand Accélérateur National d'Ions Lourds, KLAUS BARTSCHAT, Drake University, OLEG VOROV, University of North Carolina, Charlotte — Here we give analytic description of the phase transitions in the rotating Bose-Einstein condensate of weakly interacting atoms in a magnetic trap [1,2]. The analytic solution allows one to classify the instabilities in the condensates which occur when the rotational speed is increased [3]. In the case of predominantly repulsive interactions, the transition corresponds to the vortex entry the condensate. The transition to the Abrikosov state has follows if the rotational speed is increased further. In the case of predominantly attractive interactions, the transition corresponds to the escape of the condensate from the trap at the critical speed. [1] O. K. Vorov, P. Van Isacker, M. S. Hussein and K. Bartschat, Phys. Rev. Lett. 95, 230406 (2005). [2] O. K. Vorov, M. S. Hussein and P. Van Isacker, Phys. Rev. Lett. 90, 200402 (2003). [3] O. K. Vorov, P. Van Isacker, M. S. Hussein and K. Bartschat, to be submitted to Nature (2007). 1supported by NSF (USA), CEA (France)

Oleg Vorov
University of North Carolina, Charlotte

Date submitted: 09 Dec 2008

Electronic form version 1.4