

Abstract Submitted
for the MAR10 Meeting of
The American Physical Society

Isolated dimers in a 2D-like quantum spin system $\text{Li}_x\text{Cu}_{2-z}\text{Zn}_z\text{O}_2$

HUNG CHANG HSU, Novel Material Development Crystal Growth Lab, HSIANG-LIN LIU, Department of Physics at the National Taiwan Normal University, F.C. CHOU, National Taiwan University Center for Condensed Matter Sciences, NOVEL MATERIAL DEVELOPMENT CRYSTAL GROWTH LAB TEAM, MICRO OPTICAL SPECTROSCOPIC LABORATORY TEAM — The existence of isolated dimers induced by the non-magnetic dopant Zn in a 2D-like quantum spin system $\text{Li}_x\text{Cu}_{2-z}\text{Zn}_z\text{O}_2$ is found. Low Zn substitution ($< 5\%$ substituted CuO_2 chain) reduces spiral ordering temperature and a finite size effect is found to support a picture of phase separated domains with aggregated dimers near Zn dopant centers at the domain boundaries. Higher Zn substitution ($> 5\%$ substituted CuO_2 chain) introduces a novel phase transition to show a cusp shape anomaly on the magnetic susceptibility. The isotropic magnetic susceptibility reduction for high Zn level at low temperature suggests an incomplete spin gap opening and the magnetic-field dependence of the new transition temperature shows a character of weak spin-phonon coupling. Moreover, susceptibility data fitting using isolated dimer plus frustrated spin chain model describes the magnetic behaviors of samples of high Zn level between 2 - 400 K well.

Hung Chang Hsu
Department of Physics at the National Taiwan Normal University

Date submitted: 13 Nov 2009

Electronic form version 1.4