

Abstract Submitted
for the MAR10 Meeting of
The American Physical Society

Nucleation Dynamics of Particles with Directional Interactions¹

TONI PEREZ, Physics Department, Lehigh University, Bethlehem, PA, AMIT CHAKRABARTI, Department of Physics, Kansas State University, Manhattan, Kansas, JAMES D. GUNTON, Physics Department, Lehigh University, Bethlehem, PA — Protein aggregation is a subject of many studies both experimentally and theoretically since many diseases are related to undesirable protein condensation. Proteins can be modeled as globular particles with highly directional interactions. A simple model that describes this kind of protein consist of patchy particles whose interactions depend on the orientation of the patches on the surface of each particle [1]. Recently the phase diagram of a six-patch model has been reported showing the existence of different ordered phases [2]. In this work, we discuss the dynamics of the nucleation and growth process of the six-patch model using Brownian dynamics simulations. In particular we study the kinetics of the process by which chains form and ultimately collapse to spherical droplets.

[1] N. Kern and D. Frenkel, *J. Chem. Phys.* 118, 9882 (2003)

[2] H. Liu, S. K. Kumar and J. F. Douglas, *Phys. Rev. Let.* 103, 018101 (2009)

¹This work is supported by grants from the NSF and G. Harold and Leila Y. Mathers Foundation.

Toni Perez
Physics Department, Lehigh University, Bethlehem, PA

Date submitted: 18 Nov 2009

Electronic form version 1.4