

Abstract Submitted
for the MAR10 Meeting of
The American Physical Society

DFT-based first-principle calculation of Nb- and W-doped anatase TiO₂: Its stoichiometry and the formation of impurity complexes¹ HIDEYUKI KAMISAKA, TAKAHIRO SUENAGA, HISAO NAKAMURA, KOICHI YAMASHITA, The University of Tokyo — The structure and electronic states of Nb-doped anatase TiO₂ (TNO), a novel transparent conducting oxide, were calculated from the first-principle. Emphasis was put on the importance of non-stoichiometry of the system, because the high electronic conductivity emerges only in samples fabricated under a strong reductive atmosphere, and O₂ gas annealing deteriorates the conductivity. The non-stoichiometric system was modeled with periodic unit cells containing several combinations of Nb dopants, oxygen vacancies, and interstitial oxygen atom. From the result, experimental measurements of Nb 3d XPS signal and the effect of O₂ gas annealing were rationalized. Some complex structures exhibit strong energy stabilization. These structures might be a constitutive part of the electronic pathway. Similar calculations were performed for W-doped anatase TiO₂, and the difference from TNO was discussed.

¹This research was supported by a Grant for the “Elements Science and Technology Project” and the Global COE Program ”Chemical Innovation” from Ministry of Education, Culture, Sports, Science, and Technology of Japan.

Hideyuki Kamisaka
The University of Tokyo

Date submitted: 04 Jan 2010

Electronic form version 1.4