

Abstract Submitted
for the MAR10 Meeting of
The American Physical Society

Quantum Phases of Atom-Molecule Mixtures of Fermionic Atoms

NICOLAS LOPEZ VALDEZ, SHAN-WEN TSAI, University of California Riverside, CHI-YONG LIN, National Dong Hwa University, Hualien, Taiwan — Cold atom experiments have realized a variety of multicomponent quantum mixtures, including Bose-Fermi atomic mixtures. Mixtures of fermionic atoms and diatomic molecules, which are boson, have also been obtained by tuning of the interactions with external fields [1]. We study many-body correlations in such a system where the molecules are weakly bound and therefore pairs of fermionic atoms easily convert into and dissociate from the bound molecule state. This exchange mediates a long-range interaction between the fermions. We consider a simple many-body Hamiltonian that includes the destruction of fermionic atom pairs to form single bosonic molecules and vice versa [2]. We employ a functional renormalization-group approach and calculate the renormalized frequency-dependent interaction vertices and fermion self-energies. We find an instability from the disordered quantum liquid phase to a BCS phase and calculate the energy scale for the transition. The unusual frequency-dependence of this mediated interaction leads to strong renormalization of the self-energy, and also affects the couplings in the BCS channel. [1] M. Greiner, C. A. Regal, J. T. Stewart, and D. S. Jin, *Phys. Rev. Lett.* **94**, 110401 (2005) [2] E. Timmermans, K. Furuya, P. W. Milonni, and A. K. Kerman, *Phys. Lett. A* **285**, 228 (2001)

Nicolas Lopez Valdez
University of California Riverside

Date submitted: 24 Nov 2009

Electronic form version 1.4