

Abstract Submitted  
for the MAR10 Meeting of  
The American Physical Society

**Decay length of edge/surface states in topological insulators**

SHUICHI MURAKAMI, Department of Physics, Tokyo Institute of Technology and PRESTO, JST — Edge states of 2D quantum spin Hall (QSH) systems or surface states of 3D QSH systems are localized near the boundary of the systems, whereas their decay length  $\ell$  may vary. We study their generic behaviors in the presentation. We first note that  $\ell^{-1}$  plays the role of the imaginary part of the wavenumber perpendicular to the edge/surface, and  $\ell = \infty$  when the edge/surface states are absorbed into the bulk bands. We calculate  $\ell$  and discuss their behaviors by using effective models. In accordance to our expectations, we can show that the  $\ell$  is shorter when the edge/surface states are far away from the points where the edge/surface states are absorbed into the bulk bands. The edge states in HgTe quantum well or the surface states in Bi<sub>2</sub>Se<sub>3</sub> have Dirac-like dispersion and longer  $\ell$ , while in Bi ultrathin films or Bi<sub>1-x</sub>Sb<sub>x</sub>,  $\ell$  is as short as the lattice constant because the edge/surface states spread almost over the Brillouin zone. In particular, in the Dirac-like bands, the minimum  $\ell$  corresponds to the inverse of the  $k$ -space size of the Dirac cone of the edge/surface states. The QSH systems with shorter  $\ell$  will be more favorable for real-space observation of edge/surface states.

Shuichi Murakami  
Department of Physics, Tokyo Institute of Technology and PRESTO, JST

Date submitted: 20 Nov 2009

Electronic form version 1.4