

Abstract Submitted
for the MAR10 Meeting of
The American Physical Society

Spectroscopy With Free Electron Lasers DAVID BERNSTEIN, MARK BURKHARDT, SLAC and Stanford University, ANDREAS SCHERZ, JOACHIM STÖHR, WILLIAM SCHLOTTER, SLAC, YVES ACREMANN, ETH Zurich, MARTIN BEYE, TORBEN BEECK, FLORIAN SORGENFREI, ANNETTE PIETZSCH, WILFRED WURTH, ALEXANDER FÖHLISCH, FLASH and University of Hamburg — We demonstrate the feasibility of near edge x-ray absorption fine structure spectroscopy on solids by means of femtosecond soft x-ray pulses from a free-electron laser (FEL). Our experiments, carried out at the FEL at Hamburg used a special sample geometry, spectrographic energy dispersion, single shot position-sensitive detection, and a data normalization procedure that eliminates the severe fluctuations of the incident intensity in space and photon energy. As an example, we recorded the 3D_1 N_{4,5} edge absorption resonance of La³⁺ ions in LaMnO₃. Our study opens the door for x-ray absorption measurements on future x-ray FEL facilities

David Bernstein
Stanford University

Date submitted: 27 Nov 2009

Electronic form version 1.4