

Abstract Submitted  
for the MAR10 Meeting of  
The American Physical Society

**Emergent “super-solitons” following an interaction strength quantum quench across a Luttinger liquid-Mott insulating phase boundary** MATTHEW FOSTER, EMIL YUZBASHYAN, Rutgers, the State University of New Jersey — Rapid progress in cold atom experiments has motivated the study of non-equilibrium many-body dynamics following a sudden deformation of the system Hamiltonian (a “quantum quench”). Here, we consider the dynamics of localized excitations produced via a quench across a quantum phase boundary separating critical Luttinger liquid and gapped Mott insulating states. Our initial liquid ground state is labeled by a Luttinger interaction parameter  $K$ , and subject to a density-inhomogeneity forming external potential. For the Mott insulator, we employ the quantum Sine Gordon model at the Luther-Emery (LE) point. We find that over a wide range of initial  $K$  values, the quench induces the production of relativistic, non-dispersive traveling density waves, which we dub “super-solitons.” The super-solitons are generated from generic antecedent localized density lumps, and appear to be a robust feature of the post-quench dynamics. An isolated exception occurs for the case of  $K = K_{LE}$ ; here, the density dynamics are generically dispersive, and depend sensitively upon the shape of the initial inhomogeneity. We show that the super-solitons do not interact, and we demonstrate that an inhomogeneous Luttinger parameter  $K$  can be used to produce super-solitons with different characteristics in the same system.

Matthew Foster  
Rutgers, the State University of New Jersey

Date submitted: 20 Nov 2009

Electronic form version 1.4