

Abstract Submitted
for the MAR10 Meeting of
The American Physical Society

Optical Polarization of Nuclei in Bulk and Nanocrystalline Diamond¹ JONATHAN KING, JEFFREY REIMER, University of California-Berkeley — We present new results on the hyperpolarization of ^{13}C nuclear spins in diamond through optically-oriented nitrogen vacancy (NV-) defects. Optical illumination of high NV- concentration diamonds at cryogenic temperatures and 9.4 Tesla results in a negative nuclear spin temperature with measured bulk-average polarization over 5%, although local polarization may be higher. The negative spin-temperature is attributed to a population inversion within the dipolar energy levels of the NV- spin ensemble. In our quantitative model, nuclei near defects equilibrate with the NV- dipolar energy reservoir and polarization is transported to the bulk material via spin diffusion. We also investigate the nuclear hyperpolarization of NV-containing diamond nanocrystals. Such materials may be useful for surface transfer of polarization to target molecules for enhanced NMR sensitivity. Additionally, we investigate the dynamics and decoherence of the hyperpolarized nuclear spin ensemble and its interaction with electronic defect spins. Such phenomena are of fundamental interest to the use of diamond for quantum information applications.

¹The authors acknowledge support from the National Science Foundation under project ECS-0608763

Jonathan King
University of California-Berkeley

Date submitted: 20 Nov 2009

Electronic form version 1.4