

Abstract Submitted
for the MAR10 Meeting of
The American Physical Society

Alignment of Hydrogen Bonded Liquid Crystalline Block Copolymers by Magnetic Fields¹ MANESH GOPINADHAN², PAWEL MAJEWSKI³, EVAN BEACH⁴, PAUL ANASTAS⁵, CHINEDUM OSUJI⁶, Yale University, OSUJI LAB TEAM, ANASTAS LAB TEAM — Hydrogen bonding between a poly(styrene-*b*-acrylic acid) backbone and an imidazole terminated biphenyl mesogen results in the formation of a side-group liquid crystalline block copolymer (LC BCP). We use a combination of FTIR, X-ray scattering and DSC to characterize the phase behavior of the PAA-LC system, which is largely dominated by the sub-stoichiometric saturation of the binding capacity of the chain. Alignment of a hierarchically ordered lamellar BCP was performed using a 5 T magnetic field at elevated temperature in the melt state and characterized by SAXS. The system exhibits a tilted smectic structure, which on alignment by the field displays scattering patterns akin to those observed in bookshelf or chevron-type structures. These results demonstrate that simple non-covalent interactions can be used to generate LC order and thus provide a convenient handle for subsequent alignment of BCP structures by magnetic fields.

¹Support from the NSF DMR-Polymers program via DMR-0847534 is gratefully acknowledged

²Chemical Engineering

³Chemical Engineering

⁴Department of Chemistry

⁵Department of Chemistry

⁶Chemical Engineering

Manesh Gopinadhan
Yale University

Date submitted: 20 Nov 2009

Electronic form version 1.4