

Abstract Submitted
for the MAR10 Meeting of
The American Physical Society

Dipole-forbidden 5d-5f Electronic Structure in Actinide Systems

JOSEPH BRADLEY, University of Washington, SUBHRA SEN GUPTA, University of British Columbia, GERALD SEIDLER, University of Washington, GEORGE SAWATZKY, University of British Columbia, STOSH KOZIMOR, STEVEN CONRADSON, KEVIN BOLAND, DAVID CLARK, Los Alamos National Lab, UNIVERSITY OF WASHINGTON PHYSICS COLLABORATION, UNIVERSITY OF BRITISH COLUMBIA COLLABORATION, LOS ALAMOS NATIONAL LAB COLLABORATION — We report the first measurements of nonresonant inelastic x-ray scattering (NIXS) from the semi-core 5*d* levels of several actinide compounds. We find that the lowest energy electronic excitations form a rich spectrum of resonances having quite large angular momentum and whose general characteristics are predominantly dependent on the valence electron configuration rather than the nominal valence of the excited species. Comparison with atomic multiplet calculations strongly supports these observations while also illustrating a new complexity to the excited-state spectrum of actinide compounds: good agreement with experiment, subject to realistic considerations about screening effects, are achieved only when one includes the configuration interaction between the 5*f* and 6*f* states.

Joseph Bradley
University of Washington

Date submitted: 20 Nov 2009

Electronic form version 1.4