

Abstract Submitted
for the MAR10 Meeting of
The American Physical Society

Thermal conductance quantization in a T-junction¹ KEIVAN ES-FARJANI, Department of ME, MIT, Cambridge, MA, NATALIO MINGO², CEA, France — In this work, we have investigated transmission of phonons through a T-junction connecting a bulk material to a nanowire. This quantity determines the interface thermal conductance of the device and is relevant in the observation of the quantization of thermal conductance. The Green's function method is used to derive the frequency-dependence of the transmission coefficient calculated within Caroli et al's formalism. To get the frequency-dependence, we first adopt a simple nearest neighbor spring model and calculate the contribution of both linear acoustic and quadratic flexural modes of the wire, as well as the effect of the dimensionality of the bulk structure. More importantly, we also investigate the effect of the smoothness of the junction on the transmission. It is found that 2D structures smoothly connected to the wire can have a finite transmission at low frequencies if their width is finite. This will lead to a quantization of thermal conductance but not necessarily in integer multiples of the quantum of thermal conductance. Finally, results for a junction between bulk graphene and a graphene nanoribbon, using realistic force constants derived from first-principles density functional theory will be shown.

¹This research was funded by NSF.

²Department of EE, U. of California, Santa Cruz, CA

Keivan Esfarjani
Department of ME, MIT, Cambridge, MA

Date submitted: 21 Nov 2009

Electronic form version 1.4