

Abstract Submitted
for the MAR11 Meeting of
The American Physical Society

Gigantic enhancement of spin Seebeck effect by phonon drag

HIROTO ADACHI, Japan Atomic Energy Agency, KEN-ICHI UCHIDA, EIJI SAITOH, Tohoku University, JUN-ICHIRO OHE, Japan Atomic Energy Agency, SABURO TAKAHASHI, Tohoku University, SADAMICHI MAEKAWA, Japan Atomic Energy Agency — We investigate both theoretically and experimentally a gigantic enhancement of the spin Seebeck effect [K. Uchida et al., *Nature* 455, 778 (2008); C. M. Jaworski et al., *Nature Mater.* 9, 898 (2010); K. Uchida et al., *Nature Mater.* 9, 894 (2010)] in a prototypical magnet $\text{LaY}_2\text{Fe}_5\text{O}_{12}$ at low temperatures. Our theoretical analysis sheds light on the important role of phonons; the spin Seebeck effect is enormously enhanced by nonequilibrium phonons that drag the low-lying spin excitations. We further argue that this scenario gives a clue to understand the observation of the spin Seebeck effect that is unaccompanied by a global spin current, and predict that the substrate condition affects the observed signal.

Hiroto Adachi
Japan Atomic Energy Agency

Date submitted: 15 Nov 2010

Electronic form version 1.4