

Abstract Submitted
for the MAR11 Meeting of
The American Physical Society

Superconductivity at 7.7 K in new hexagonal bronze Hg_xReO_3

KENYA OHGUSHI, University of Tokyo, JST-TRIP, AYAKO YAMAMOTO, RIKEN, JST-TRIP, YOKO KIUCHI, CHANDREYEE GANGULI, University of Tokyo, KAZUYUKI MATSUBAYASHI, YOSHIYA UWATOKO, University of Tokyo, JST-TRIP, HIDENORI TAKAGI, RIKEN, University of Tokyo, JST-TRIP — We have successfully synthesized a new rhenium-based hexagonal bronze material, Hg_xReO_3 , which exhibits superconductivity with the transition temperature $T_c = 7.7$ K at ambient pressure and 11.1 K at 4 GPa. This compound is a superconductor with the highest T_c among hexagonal bronzes. Moreover, it presents the novel crystallographic feature that $(\text{Hg}_2)^{2+}$ polycations, in contrast to monatomic cations in known hexagonal bronzes, are incorporated into open channels. There is evidence that conducting electrons tightly couple with Hg-related phonons. Our results inspire detailed studies on the role of the rattling phonon in the occurrence of superconductivity in the hexagonal bronzes.

Kenya Ohgushi
University of Tokyo, JST-TRIP

Date submitted: 16 Nov 2010

Electronic form version 1.4