

Abstract Submitted
for the MAR11 Meeting of
The American Physical Society

Two-Dimensional quantum dynamic in a dc SQUID¹ FLORENT LECOCQ, Institut Neel, CNRS, Grenoble, France, I.M. POP, Z. PENG, I. MATEI, C. NAUD, F.W. HEKKING, W. GUICHARD, O. BUISSON, Institut Neel and LPMMC, CNRS, Grenoble, France, R. DOLATA, A.B. ZORIN, PTB, Braunschweig, Germany — The dynamics of a dc SQUID presents a large variety of quantum effects at very low temperature such as 2D MQT signature, multilevel and phase qubit dynamics. We have shown that along the zero current bias line, the quantum dynamics is protected from current fluctuations. Along this line, the potential is quadratic-quartic and enhanced phase qubit properties have been demonstrated² When the dc SQUID loop inductance is of the order of the Josephson inductance the dynamic becomes two dimensional. As a consequence, in addition to the oscillation mode producing the phase qubit, a second oscillation mode exists, called transverse mode. Here we report spectroscopic evidence and coherence properties of both oscillators as well as coherent oscillations between the quantum states of these two coupled oscillators.

¹Supported by the EU project EuroSQIP and SOLID, and ANR QUANTJO.

²E. Hoskinson et al, Phys. Rev. Lett. 102, 097004 (2009)

Florent Lecocq
Institut Neel, CNRS, Grenoble, France

Date submitted: 21 Nov 2010

Electronic form version 1.4