

Abstract Submitted
for the MAR11 Meeting of
The American Physical Society

London penetration depth of λ -(BETS) $_2$ GaCl $_4$ superconductor

JASON MURPHY, H. KIM, K. CHO, M.A. TANTAR, Iowa State University/Ames Laboratory, A. KOBAYASHI, University of Tokyo, Japan, H. KOBAYASHI, Department of Chemistry, Nihon University, Japan, R. PROZOROV, Iowa State University/Ames Laboratory — After many years of study, it is still unclear whether the superconductivity of organic charge transfer salts is conventional or not. In particular, symmetry of the superconducting gap is an important unresolved question. The gap anisotropy and nodal structure can be probed via precision measurements of London penetration depth, $\lambda(T)$, for which tunnel-diode resonator has been developed over past 20 years. Here we report $\lambda(T)$ in λ -(BETS) $_2$ GaCl $_4$ crystals ($T_c \sim 5$ K) measured down to 400 mK. We find that different samples exhibit different widths of superconducting transition, ΔT_c . It is known for organic superconductors that cooling rate from room temperature is important for the properties probed at the low temperatures. We discuss observed variation of ΔT_c with the cooling rate as well as sample size and aspect ratio. Results are compared with thermal conductivity measurements [1].

[1] M. A. Tanatar, T. Ishiguro, H. Tanaka, H. Kobayashi, Phys.Rev. B **66**, 134503 (2002).

Jason Murphy
Iowa State University/Ames Laboratory

Date submitted: 21 Nov 2010

Electronic form version 1.4