## Abstract Submitted for the MAR11 Meeting of The American Physical Society

Observation of double hysteresis in a MnMn<sub>6</sub>(CH<sub>3</sub>H<sub>5</sub>O<sub>3</sub>)<sub>3</sub> singlemolecule magnet KLAUS GIEB, WOLFGANG KROENER, PAUL MULLER, Department of Physics and Interdisciplinary Center for Molecular Materials (ICMM), Universitaet Erlangen-Nuernberg, Germany, CARL-GEORG FREIHERR VON RICHTHOFEN, THORSTEN GLASER, Fakultaet fuer Chemie - Lehrstuhl Anorganische Chemie I, Universitaet Bielefeld, Germany — We report on high field and low temperature magnetization measurements of a novel MnMn<sub>6</sub>(CH<sub>3</sub>H<sub>5</sub>O<sub>3</sub>)<sub>3</sub> complex. A home-made micro-Hall-probe magnetometer was used to perform the characterization at mK temperatures and fields up to 17 T. Most 3d-ion based singlemolecule magnets, known up to now, have a spin ground state well separated from the first excited state, leading to the formation of giant spin at low temperatures. In contrast to this situation, the ground state (S=6) of the present complex can already be exited at moderate magnetic fields. Surprisingly, magnetic hysteresis was observed for both the ground state and the first exited state leading to a double hysteresis in the low temperature magnetization measurements. The blocking temperature was found to be  $T_B \approx 1.3 \,\mathrm{K}$ . Origin and possible consequences of this unusual behavior will be discussed.

> Klaus Gieb Uni Erlangen

Date submitted: 22 Nov 2010 Electronic form version 1.4