

Abstract Submitted  
for the MAR11 Meeting of  
The American Physical Society

**Scanning Tunneling Microscopy Study of Quantum Cobalt Chains**<sup>1</sup>

NADER ZAKI, Columbia University, DANDA ACHARYA, Pacific Northwest National Laboratory, DENIS POTAPENKO, Columbia University, PETER JOHNSON, PERCY ZAHL, PETER SUTTER, Brookhaven National Laboratory, RICHARD OSGOOD, Columbia University — We recently reported [1] on a new surface phase of the Co-vicinal-Cu(111) system, which exhibits self-assembled uniform Co quantum wires that are stable at 300K. STM images show that the wires form along the leading edge of the step rise, differentiating it from previously theoretically predicted atomic-wire phases as well as experimentally observed step-island formation. Our observations allow us to comment on the formation kinetics of the atomic-wire phase and on the fit of our data to a recently developed lattice-gas model. LT-STM measurements, taken on self-assembled Co chains, reveal a charge-density modulation that is dependent on tip bias. These charge-modulations are observed for tip-bias relatively far away from the Fermi level, both at negative and positive bias. We present arguments for the identification of these modulations as due to either charge-density-like waves (CDW) or excited states of this 1-D system.

[1] N. Zaki et al, Phys. Rev. B 80, 155419 (2009)

<sup>1</sup>Department of Energy Contract No. DE-FG 02-04-ER-46157

Nader Zaki  
Columbia University

Date submitted: 20 Dec 2010

Electronic form version 1.4