

Abstract Submitted
for the MAR11 Meeting of
The American Physical Society

**Rotational Diffusion of Plasmon-Resonant Gold Nanorods for
Depth-Resolved Microrheology Using Optical Coherence Tomography¹**

AMY OLDENBURG, RAGHAV CHHETRI, University of North Carolina at Chapel Hill, KRYSYTIAN KOZEK, AARON JOHNSTON-PECK, JOSEPH TRACY, North Carolina State University — The ability to perform microrheology in optically thick samples would enable analysis of bulk tissues. Optical coherence tomography (OCT) provides imaging several mean free scattering path lengths into tissue. In this study we report the use of plasmon-resonant gold nanorods as microrheological sensors in OCT. Nanorods exhibit a longitudinal mode that is excited when they are oriented parallel to the polarization of the incident light, which is favorable for passive microrheology using polarized light to monitor their rotational diffusion. We demonstrate measurements of the rotational diffusion of unconfined, colloidal gold nanorods using polarization-sensitive OCT, and validate the Stokes-Einstein relationship for the nanorods in simple fluids of varying viscosity. We then show that OCT provides depth-resolved imaging of fluid viscosity through measurements of the rotational diffusion rate of the nanorods.

¹We acknowledge support from the Carolina Cancer Center for Nanotechnology Excellence (C-CCNE NIH (NCI) #U54CA119343).

Amy Oldenburg
University of North Carolina at Chapel Hill

Date submitted: 23 Nov 2010

Electronic form version 1.4