

Abstract Submitted  
for the MAR11 Meeting of  
The American Physical Society

**Doping dependence of the optical spectra in  $\text{Fe}_{1.02}\text{Te}_{1-x}\text{Se}_x$** <sup>1</sup>  
CATALIN MARTIN, X. XI, K. MILLER, D. KOUKIS, University of Florida, J. HU, Z.Q. MAO, Tulane University, G.L. CARR, Brookhaven National Laboratory, D.B. TANNER, University of Florida — We measured the optical reflectivity and extracted the optical conductivity over a broad spectral range of the Fe-chalcogenide  $\text{Fe}_{1.02}\text{Te}_{1-x}\text{Se}_x$ , at various doping levels, from  $x=0.12$  to  $x=0.45$ . Except for the highest concentration of Se ( $x=0.45$ ), optical properties show unusual frequency and temperature dependence. The zero frequency Drude peak is absent in optical conductivity, a plasma edge cannot be unambiguously defined and both reflectivity and optical conductivity have non-monothonic temperature dependence. For the superconducting samples, we found a weak enhancement of the reflectivity in the far-infrared region upon cooling below  $T_c$  and its possible association with the superconducting gap will be discussed.

<sup>1</sup>Supported by the US DOE through contracts DE-FG02-02ER45984 and DE-AC02-98CH10886.

Catalin Martin  
University of Florida

Date submitted: 22 Dec 2010

Electronic form version 1.4