

Abstract Submitted
for the MAR11 Meeting of
The American Physical Society

Dynamics of endohedral hydrogen in C₆₀: Infrared study¹ MIN GE, U. NAGEL, D. HUVONEN, T. ROOM, Nat. Inst. of Chem. Phys. Biophys., Tallinn, Estonia, S. MAMONE, M.H. LEVITT, M. CARRAVETTA, Southampton Uni., UK, Y. MURATA, K. KOMATSU, Kyoto Uni., Japan, J.Y.-C. CHEN, N.J. TURRO, Columbia Uni., NY10027 — Infrared spectra of endohedral hydrogen isotopomers H₂, D₂, and HD were measured in the temperature range from 6 to 300 K. A model of a vibrating rotor in a spherical potential together with the translational motion induced dipole moment theory was used to explain the positions and intensities of IR absorption lines. By measuring spectra of a *para* enriched sample of H₂@C₆₀, we confirmed the assignment of lines to *ortho*- and *para*-H₂. Inside C₆₀ cage the rotation of hydrogen is unhindered and the translation is quantized and coupled to the rotation. The isotropic and translation-rotation coupling part of the potential are anharmonic and depend on the vibrational states of hydrogen. The analysis of the isotopic effect on the IR transition intensities and interaction potential in the C₆₀ cage is presented.

¹Support by the Estonian Ministry of Education and Research grant SF0690029s0 and SF grants ETF7011, ETF8170 and JD187 is acknowledged.

Toomas Room
Nat. Inst. of Chem. Phys. Biophys., Tallinn, Estonia

Date submitted: 23 Dec 2010

Electronic form version 1.4