

Abstract Submitted
for the MAR11 Meeting of
The American Physical Society

Highly confined 2D electron gas in SrTiO₃/SrRuO₃ superlattices¹

MARCOS VERÍSSIMO-ALVES, CITIMAC, Universidad de Cantabria, Spain, DANIEL BILC, Universite de Liège, Belgium, PABLO GARCÍA-FERNÁNDEZ, CITIMAC, Universidad de Cantabria, Spain, PHILIPPE GHOSEZ, Universite de Liège, Belgium, JAVIER JUNQUERA, CITIMAC, Universidad de Cantabria, Spain — In recent years, experimental advances in the creation of atomically smooth interfaces between complex oxides have made possible the observation of unusual electronic phenomena such as the formation of 2D electron gases confined to the interface between insulating compounds. In this work, we explore an alternative for creating a 2D electron gas in oxide interfaces by means of a (SrTiO₃)₅/(SrRuO₃)₁ superlattice. LSDA+U and hybrid functional B1-WC DFT calculations with full relaxation of atomic geometry show the presence of a 2D electron gas, and the system can be described with $U = 4$ eV. Electronic correlations thus become important with quantum confinement: the system is a minority spin half metal with a magnetic moment of $\mu = 4\mu_B$. Degenerate Ru 4d_(xz,yz) states are the main responsible for the DOS at the Fermi level, whose shape strongly resembles that of 1D free-electron gases. Our results suggest that the material could find uses in spintronic devices.

¹We thank financial support from FIS2009-12721-C04-02 and CP-FP 228989-2 Projects.

Javier Junquera
CITIMAC, Universidad de Cantabria, Spain

Date submitted: 27 Dec 2010

Electronic form version 1.4