

Abstract Submitted
for the MAR11 Meeting of
The American Physical Society

Large-Scale Molecular Dynamics Simulations of Shock-Induced Plasticity in Tantalum Single Crystals¹ R. RAVELO², University of Texas-El Paso, QI AN, California Institute of Technology, T.C. GERMANN, B.L. HOLIAN, Los Alamos National Laboratory — We report on large-scale non-equilibrium molecular dynamics (NEMD) simulations of shock wave compression in Ta single crystals. The atomic interactions are modeled via a recently developed and optimized embedded-atom method (EAM) potential for Ta, which reproduces the equation of state up to 200 GPa. We examined the elastic-plastic transition and shock wave structure for wave propagation along the low index directions: (100), (110) and (111). Shock waves along (100) and (111) exhibit an elastic precursor followed by a plastic wave for particle velocities below 1.1 km/s for (100) and 1.4 km/s for (111). The nature of the plastic deformation along (110) is dominated by twinning for pressures above 40 GPa.

¹Part of this work was supported by the U.S. Department of Energy under contract DE-AC5206NA25396.

²Computational Physics Division, Los Alamos National Laboratory

Ramon Ravelo
University of Texas-El Paso

Date submitted: 27 Nov 2010

Electronic form version 1.4