

Abstract Submitted
for the MAR11 Meeting of
The American Physical Society

Origin of the Giant Honeycomb Network of Quinones on Cu(111)¹

T.L. EINSTEIN, KWANGMOO KIM, U. Maryland, JON WYRICK, ZHIHAI CHENG, LUDWIG BARTELS, UC Riverside, KRISTIAN BERLAND, PER HYLDGAARD, Chalmers U. Tech. — We discuss the factors that lead to the amazing regular giant honeycomb network formed by quinones on Cu(111). Using a related lattice gas model with many characteristic energies, we can reproduce many experimental features. These models require a long-range attraction, which can be attributed to indirect interactions mediated by the Shockley surface state of Cu(111). However, Wyrick's preceding talk gave evidence that the network self-selects for the size of the pore rather than for the periodicity of the superstructure, suggesting that confined states are the key ingredient. We discuss this phenomenon in terms of the magic numbers of 2D quantum dots. We also report calculations of the effects of anthraquinones (AQ) in modifying the surface states by considering a superlattice of AQ chains with various separations. We discuss implications of these results for tuning the electronic states and, thence, superstructures.

¹Supported by (TLE) NSF CHE 07-50334 & UMD MRSEC DMR 05-20471, (JW & LB) NSF CHE NSF CHE 07-49949, (KB & PH) Swedish Vetenskapsrådet VR 621-2008-4346.

Theodore Einstein
U. Maryland, College Park

Date submitted: 27 Nov 2010

Electronic form version 1.4