

Abstract Submitted
for the MAR11 Meeting of
The American Physical Society

The Effect of Magnetic Anisotropy on Colossal Electroresistance in Manganites¹ ALESSANDRA GALLASTEGUI, RAFIYA JAVED, HYOUNG-JEEN JEEN, AMLAN BISWAS, Department of Physics, University of Florida, Gainesville, FL 32611 — The combined effect of long range strain interactions and disorder on a first order transition leads to micrometer scale phase separation in hole-doped manganese oxides (manganites). The coexisting phases are ferromagnetic metallic (FMM), charge ordered insulating (COI), and paramagnetic insulating (PMI) and at certain temperatures these phases behave like a fluid under the influence of magnetic and electric fields. We will present magnetotransport data on nano/micro-structures of the manganite $(\text{La}_{1-y}\text{Pr}_y)_{0.67}\text{Ca}_{0.33}\text{MnO}_3$ (LPCMO) which show that the FMM phase behaves like a fluid in an electric field. In fact, due to the magnetic anisotropy of our materials, the behavior of the coexisting phases is reminiscent of a ferrofluid.

¹NSF DMR-0804452

Alessandra Gallastegui
Department of Physics, University of Florida, Gainesville, FL 32611

Date submitted: 19 Nov 2010

Electronic form version 1.4