

Abstract Submitted
for the MAR11 Meeting of
The American Physical Society

Burnett-Cattaneo Continuum Theory for Shock Waves¹ B.L. HOLIAN, Los Alamos National Laboratory, M. MARESCHAL, Universite Libre de Bruxelles, R. RAVELO, University of Texas-El Paso — We model strong shockwave propagation, both in the ideal gas and in the dense Lennard-Jones fluid, using a refinement of earlier work ², which accounts for the cold compression by a nonlinear, Burnett-like, strain-rate dependence of the thermal conductivity, and relaxation of temperature components on the hot, compressed side of the shock front. The relaxation of the disequilibrium among the three components of the kinetic temperature, namely, the difference between the temperature in the direction of a planar shock wave and those in the transverse directions, particularly in the region near the shock front, is accomplished by a rigorous application of the Cattaneo-Maxwell relaxation equation to a reference state, namely, the steady shockwave solution of linear Navier-Stokes-Fourier theory, along with the nonlinear Burnett heat-flux term. Our new continuum theory is in nearly quantitative agreement with non-equilibrium molecular-dynamics simulations under strong shockwave conditions.

¹Part of this work supported by the U.S. Department of Energy under contract DE-AC5206NA25396.

²B.L. Holian, M. Mareschal, and R. Ravelo, *J. Chem. Phys.* **133**, 114502 (2010)

Ramon Ravelo
University of Texas-El Paso

Date submitted: 28 Nov 2010

Electronic form version 1.4