

Abstract Submitted  
for the MAR11 Meeting of  
The American Physical Society

**Raman Maps of Carbon Nanoscrolls and their Bundles**

GUANGJUN CHENG, IRENE CALIZO, XUELEI LIANG, BRENT SPERLING, JAMES MASLAR, CURT A. RICHTER, ANGELA R. HIGHT WALKER, National Institute of Standards and Technology (NIST), Gaithersburg, MD 20899 — Recent theoretical simulations and experimental data have shown that carbon nanoscrolls, formed by rolling a single layer of graphene, exhibit different electronic and optical properties from carbon nanotubes or the planar graphene sheet. They hold promise in a variety of applications areas such as energy storage and nanomechanics. Here, we present our investigation on the formation of carbon nanoscrolls and their bundles from large-area graphene originally grown on copper foil by chemical vapor deposition. When graphene is transferred from a copper foil onto a silicon wafer using wet chemistry, both filled and unfilled carbon nanoscrolls are produced upon the rupture of graphene. These nanoscrolls can further form bundles. The results from Raman mapping, optical microscopy, scanning electron microscopy, and atomic force microscopy measurements will be presented and the formation mechanism will be discussed.

Angela R. Hight Walker  
National Institute of Standards and Technology (NIST),  
Gaithersburg, MD 20899

Date submitted: 28 Nov 2010

Electronic form version 1.4