

Abstract Submitted
for the MAR11 Meeting of
The American Physical Society

Terahertz and Infrared Conductivity of Large-Area Graphene

LEI REN, QI ZHANG, TAKASHI ARIKAWA, LAYLA G. BOOSHEHRI, JUNICHIRO KONO, Department of Electrical and Computer Engineering, Rice University, ZHONG JIN, ZHENGZONG SUN, ZHENG YAN, JAMES M. TOUR, Department of Chemistry, Rice University — Graphene is predicted to offer new opportunities for terahertz (THz) science and technology. Its zero-gap linear band dispersion is expected to lead to exotic nonlinear electromagnetic properties, which can be probed through frequency-dependent conductivity measurements. Here, we use THz time-domain spectroscopy and Fourier-transform infrared spectroscopy to investigate carrier dynamics in large-area graphene grown by chemical vapor deposition. We studied both nitrogen-doped and nominally-undoped graphene; the latter had accidental doping presumably through air and acid exposure. Absorption increased with the number of graphene layers and was larger in the nominally-undoped samples especially in the 0.2-2.2 THz range. For the highest-mobility samples, we observed Drude-like frequency dependence in the THz range. Further measurements in a wider spectral range are in progress to understand the differences between these samples and the interplay between intra-band and inter-band dynamics.

Lei Ren
Department of Electrical and Computer Engineering, Rice University

Date submitted: 19 Nov 2010

Electronic form version 1.4