

Abstract Submitted  
for the MAR11 Meeting of  
The American Physical Society

**Effect of doping on the crystalline structure and superconductivity properties in  $\text{Ca}_{1-x}\text{Na}_x\text{Fe}_2\text{As}_2$  single crystals**<sup>1</sup> LEONARDO CIVALE, NESTOR HABERKORN, BORIS MAIOROV, MARCELO JAIME, Los Alamos National Laboratory, Los Alamos, NM, USA, G. CHEN, W. YU, Renmin University of China, Beijing, China — We have studied the crystalline structure and the superconducting properties of  $\text{Ca}_{1-x}\text{Na}_x\text{Fe}_2\text{As}_2$  single crystals for various levels of the chemical doping ( $x$ ). We performed a comparative analysis of the angular dependent  $H_{c2}(\Theta)$  (where  $\Theta$  is the angle between the magnetic field and the  $c$  axis) for  $x \sim 0.5$  and  $x \sim 0.75$ , corresponding to  $T_c \sim 19$  K and 33 K, respectively. We found that in both cases  $H_{c2}(\Theta)$  near  $T_c$  exhibits a single band character with the same anisotropy  $\gamma \sim 1.8$ . In the crystal with  $T_c \sim 33$  K we detected the presence of a narrow vortex-liquid phase, in agreement with the expectation from estimates based on the Lindemann criterion and the Ginzburg number. We also found large and anisotropic flux creep rates, with temperature dependences that indicate glassy relaxation. We analyzed those results in terms of single vortex and collective pinning regimes associated with random and correlated disorder.

<sup>1</sup>Research supported by the U.S. Department of Energy, Office of Basic Energy Sciences, Division of Materials Sciences and Engineering.

Leonardo Civale  
Los Alamos National Laboratory, Los Alamos, NM, USA

Date submitted: 08 Dec 2010

Electronic form version 1.4