

Abstract Submitted
for the MAR11 Meeting of
The American Physical Society

The Spin-Lattice Relaxation of Hyperpolarized ^{89}Y Complexes

ASHISH JINDAL, LLOYD LUMATA, UT Southwestern Med. Cntr., YIXUN XING, UT Dallas, MATTHEW MERRITT, UT Southwestern Med. Cntr., PIYU ZHAO, UT Dallas, CRAIG MALLOY, DEAN SHERRY, ZOLTAN KOVACS, UT Southwestern Med. Cntr. — The low sensitivity of NMR can be overcome by dynamic nuclear polarization (DNP). However, a limitation to the use of hyperpolarized materials is the signal decay due to T_1 relaxation. Among NMR-active nuclei, ^{89}Y is potentially valuable in medical imaging because in chelated form, pH-sensitive agents can be developed. ^{89}Y also offers many attractive features – 100 % abundance, a 1/2 spin, and a long T_1 , up to 10 min. Yet, developing new ^{89}Y complexes with even longer T_1 values is desirable. Designing such complexes relies upon understanding the mechanism(s) responsible for T_1 relaxation. We report an approach to hyperpolarized T_1 measurements that enabled an analysis of relaxation mechanisms by selective deuteration of the ligand backbone, the solvent or both. Hyperpolarized ^{89}Y – DTPA, DOTA, EDTA, and deuterated EDTA complexes were studied. Results suggest that substitution of low-gamma nuclei on the ligand backbone as opposed to that of the solvent most effectively increase the ^{89}Y T_1 . These results are encouraging for in vivo applications as the presence of bound water may not dramatically affect the T_1 .

Ashish Jindal
UT Southwestern Med. Cntr.

Date submitted: 19 Nov 2010

Electronic form version 1.4