

Abstract Submitted  
for the MAR11 Meeting of  
The American Physical Society

**Predicting Sommerfeld coefficients for heavy-fermion materials**

MUNEHISA MATSUMOTO, SERGEY SAVRASOV, UC Davis, JUNYA OTSUKI, Tohoku University, Japan — From electronic-structure calculation standpoint we predict the appearance of heavy-fermion behavior for Cerium and Plutonium-based materials. Local-density approximation (LDA) combined with dynamical-mean field theory (DMFT) formulated for localized f-electrons gives an efficient realistic Kondo-lattice description [1] for the target materials, yielding a quasiparticle renormalization factor  $z_c$  for conduction electrons. We invert the data to get the quasiparticle renormalization factor  $z_f$  for f-electrons, and restore the effective total density of states to predict the Sommerfeld coefficient  $\gamma$ . Summarizing our data on an analogous “Doniach phase diagram” plotted for  $z_c$ ,  $z_f$ , and  $\gamma$ ,  $\gamma$  is found to have a peak around the magnetic quantum critical point and Ce-115’s are found to have the largest  $\gamma$ ’s among target materials.

[1] PRL **103**, 096403 (2009).

Munehisa Matsumoto  
UC Davis

Date submitted: 07 Dec 2010

Electronic form version 1.4