

Abstract Submitted
for the MAR12 Meeting of
The American Physical Society

**Morphology and Dynamics of Liquid Crystalline
Molecules Confined in Nano-pores**

MALGORZATA JASIURKOWSKA, WILHELM KOSSACK, ROXANA ENE, CIPRIAN IACOB, WYCLIFFE KIPNUSU, PERIKLIS PAPADOPOULOS, JOSHUA SANGORO, University of Leipzig, Germany, MARIA MASSALSKA-ARODZ, Polish Academy of Sciences, Krakow, Poland, FRIEDRICH KREMER, University of Leipzig, Germany, UNIVERSITY OF LEIPZIG, GERMANY TEAM, POLISH ACADEMY OF SCIENCES, KRAKOW, POLAND COLLABORATION — Broadband dielectric and Infrared spectroscopy are combined to study the molecular dynamics of the liquid crystalline compounds belong isothiocyanatobiphenyl homologous series (abbreviated as nBT) confined in pores of mean diameters from 4 nm to 10.5 nm. In bulk, the studied substances show only one liquid crystalline phase: the SmE phase with orthorhombic arrangement within the molecular layers. In contrast to well-known bulk dielectric properties of nBTs, confinement leads to modification of the molecular dynamics. Two relaxation processes are detected. The slower process corresponds to molecular reorientation around short axis and it is faster in pores than in bulk. The second process is attributed to a librational motion of the molecules close to the walls. Both processes exhibit an Arrhenius-type temperature dependence. Detailed analysis of the temperature dependent infrared spectra indicates the different impact of confinement on the rigid and flexible molecular units of nBTs. Transition Moment Orientational Analysis is employed to explore the orientational order of molecules in pores.

Malgorzata Jasiurkowska
Institute of Experimental Physics I, University of Leipzig,
04103 Leipzig, Germany

Date submitted: 26 Nov 2011

Electronic form version 1.4