

Abstract Submitted
for the MAR12 Meeting of
The American Physical Society

First principles computer simulations of $\text{Li}_{10}\text{GeP}_2\text{S}_{12}$ and related lithium superionic conductors¹ N.A.W. HOLZWARTH, Wake Forest University — A recent paper by Kamaya *et al.*² reported a new crystalline superionic conductor having a compact tetrahedral structure and a stoichiometry of $\text{Li}_{10}\text{GeP}_2\text{S}_{12}$. The room temperature conductivity was reported to be 0.01 S/cm, comparable to liquid electrolyte conductivities and five times higher the compositionally related thio-LISICON material $\text{Li}_{3.25}\text{Ge}_{0.25}\text{P}_{0.75}\text{S}_4$ developed earlier.³ This talk will present a progress report on our work to perform first principles computer simulations for these materials focussing on the structural, stability, and Li ion mobility properties of idealized crystalline models. From the perspective of our previous studies of Li ion conductivity in lithium thiophosphate electrolytes,⁴ the effects of introducing Ge can be assessed.

¹Supported by NSF Grants DMR-0705239 and DMR-1105485.

²N. Kamaya *et al.*, *Nature Materials* **10**, 682 (2011).

³R. Kanno *et al.*, *J. Electrochem. Soc.* **148**, A742 (2001).

⁴N. A. W. Holzwarth *et al.*, *J. Power Sources* **196**, 6870 (2011).

N. A. W. Holzwarth
Wake Forest University

Date submitted: 15 Dec 2011

Electronic form version 1.4