

Abstract Submitted  
for the MAR12 Meeting of  
The American Physical Society

**A Comparison of Different Treatments of Spin-Orbit  
Coupling within the GW Approach<sup>1</sup>**

BRAD BARKER, University of California-Berkeley, Lawrence Berkeley National Laboratory, JACK DESLIPPE, Lawrence Berkeley National Laboratory, OLEG V. YAZYEV, Ecole Polytechnique Federale de Lausanne (EPFL), Switzerland, STEVEN G. LOUIE, University of California-Berkeley, Lawrence Berkeley National Laboratory — Spin-orbit coupling is responsible for the unique electronic properties of many fascinating materials such as topological insulators. We have developed and implemented an approach in which the effects of spin-orbit interactions to the quasiparticle band structure are incorporated within the GW approach, employing spinor wavefunctions computed at the DFT level with fully relativistic pseudopotentials. We compare these results to separate calculations where spin-orbit coupling is applied as a perturbation either before or after the GW calculation of the band structure. We apply these methods to the properties of materials with heavy ion cores to determine possible differences from the different schemes.

<sup>1</sup>This work was supported by National Science Foundation Grant No. DMR10-1006184, the U.S. Department of Energy under Contract No. DE-AC02-05CH11231. Computational resources have been provided by DOE at Lawrence Berkeley National Laboratory's NERSC facility

Brad Barker  
University of California-Berkeley, Lawrence Berkeley National Laboratory

Date submitted: 11 Nov 2011

Electronic form version 1.4