

Abstract Submitted  
for the MAR12 Meeting of  
The American Physical Society

**Topological States and Adiabatic Pumping in Quasicrystals<sup>1</sup>**

YAAKOV KRAUS, YOAV LAHINI, ZOHAR RINGEL, MOR VERBIN, ODED ZILBERBERG, Weizmann Institute of Science — We find a connection between quasicrystals and topological matter, namely that quasicrystals exhibit non-trivial topological phases attributed to dimensions higher than their own [1]. Quasicrystals are materials which are neither ordered nor disordered, i.e. they exhibit only long-range order [2]. This long-range order is usually expressed as a projection from a higher dimensional ordered system. Recently, the unrelated discovery of Topological Insulators [3] defined a new type of materials classified by their topology. We show theoretically and experimentally using photonic lattices, that one-dimensional quasicrystals exhibit topologically-protected boundary states equivalent to the edge states of the two-dimensional Integer Quantum Hall Effect. We harness this property to adiabatically pump light across the quasicrystal, and generalize our results to higher dimensional systems. Hence, quasicrystals offer a new platform for the study of topological phases while their topology may better explain their surface properties.

[1] Y. E. Kraus, Y. Lahini, Z. Ringel, M. Verbin, and O. Zilberberg, arXiv:1109.5983 (2011).

[2] C. Janot, *Quasicrystals* (Clarendon, Oxford, 1994), 2nd ed.

[3] M. Z. Hasan and C. L. Kane, Rev. Mod. Phys. **82**, 3045 (2010).

<sup>1</sup>Authors appear in alphabetical order.

Oded Zilberberg  
Weizmann Institute of Science

Date submitted: 19 Nov 2011

Electronic form version 1.4