

Abstract Submitted
for the MAR12 Meeting of
The American Physical Society

**A Cahn-Hilliard model of vascularized tumor growth
in a complex evolving confinement using a diffuse domain
approach¹**

YAO-LI CHUANG, University of New Mexico, Dept. Pathology, VITTORIO CRISTINI, University of New Mexico, Dept. Pathology and Dept. Chemical Engineering, YING CHEN, XIANG-RONG LI, University of California, Irvine, Dept. Math, HERMANN FRIEBOES, University of Louisville, Dept. Bioengineering, JOHN LOWENGRUB, University of California, Irvine, Dept. Math — Understanding the spatiotemporal evolution of tumor growth is essential for developing effective strategies to treat cancers. Various studies have suggested that spatial heterogeneity during tumors growth is a key factor associated with subsequent tumor invasion and the effectiveness of chemotherapy. Spatial heterogeneity may arise due to morphological instability of the tumors and the complex tissue structure surrounding the tumors. In previous works, we have used a Cahn-Hilliard tumor growth model to study the morphological instability for tumors in non-resisting tissues. However, most tumors are surrounded by complex tissue structures and confined in the capsules of some organs or between certain basement membranes. The capsules and basement membranes may be distorted by interacting with the evolving tumors, affecting the morphological instability. Here we adopt a novel diffuse domain approach to adapt our previous Cahn-Hilliard model for tumor growth in such complex evolving environments. As an example, we apply the model to simulate the evolution of lymphoma in a lymph node, incorporating also the tumor-induced angiogenesis.

Yao-li Chuang

University of New Mexico, Dept. Pathology

¹supported by NIH-PSOC grant 1U54CA143907-01

Date submitted: 11 Nov 2011

Electronic form version 1.4