

Abstract Submitted
for the MAR12 Meeting of
The American Physical Society

Quantum pumping of electrons at the Josephson frequency

FRANCESCO GIAZOTTO, PANAYOTIS SPATHIS, STEFANO RODDARO, SUBHAJIT BISWAS, FABIO TADDEI, NEST, Istituto Nanoscienze-CNR and Scuola Normale Superiore, Piazza S. Silvestro 12, I-56127 Pisa, Italy, MICHELE GOVERNALE, School of Chemical and Physical Sciences and MacDiarmid Institute for Advanced Materials and Nanotechnology, Victoria University of Wellington, P.O. B, LUCIA SORBA, NEST, Istituto Nanoscienze-CNR and Scuola Normale Superiore, Piazza S. Silvestro 12, I-56127 Pisa, Italy — A macroscopic fluid pump works according to the law of Newtonian mechanics and transfers a large number of molecules per cycle. By contrast, a nano-scale charge pump can be thought as the ultimate miniaturization of a pump, with its operation being subject to quantum mechanics and with only few electrons or even fractions of electrons transferred per cycle. It generates a direct current in the absence of an applied voltage exploiting the time-dependence of some properties of a nano-scale conductor. So far, nano-scale pumps have been realised only in system exhibiting strong Coulombic effects, whereas evidence for pumping in the absence of Coulomb-blockade has been elusive. Here we report the experimental detection of charge flow in an unbiased InAs nanowire embedded in a superconducting quantum interference device (SQUID). In this system, pumping occurs via the cyclic modulation of the phase of the order parameter of different superconductors. The symmetry of the current with respect to the enclosed flux and SQUID current is a discriminating signature of pumping. Currents exceeding 20 pA are measured at 250 mK, and exhibit symmetries compatible with a pumping mechanism.

Francesco Giazotto
NEST, Istituto Nanoscienze-CNR and Scuola Normale Superiore,
Piazza S. Silvestro 12, I-56127 Pisa, Italy

Date submitted: 13 Dec 2011

Electronic form version 1.4