

Abstract Submitted
for the MAR13 Meeting of
The American Physical Society

Closed Channel Amplitude in a Many Body Feshbach System

NICOLAS LOPEZ, University of California Riverside, EDDY TIMMERMANS, Los Alamos National Lab, SHAN-WEN TSAI, University of California Riverside — Near a narrow Feshbach resonance (with magnetic field width 10 mG or smaller) the ultra-cold atom interactions acquire an effective range that can be comparable to the average inter-particle distance. Although requiring a more accurate magnetic field control than their broad counterparts, the narrow Feshbach resonances can free cold atom physics from its straightjacket of the contact interaction paradigm. The finite-range effects can give rise to roton features in the phonon dispersion of dilute Bose-Einstein condensates (BEC's) and BEC's can support a ground state with modulated density patterns that breaks translational symmetry. We show that the finite range interaction is the consequence of the time-delay in atom-atom collisions. The narrow regime is also the parameter region in which the interacting atoms can spend a significant fraction of their time in the spin-rearranged (also called “closed”) channel. To study the interaction physics we describe two atoms in a harmonic trap, interacting near a narrow resonance. We find the fraction of time that the atoms spend in the closed channel at fixed magnetic field and we extend this result to estimate the fraction of time that a distinguishable atom moving through a BEC spends in the closed channel state, quasibound to BEC-atoms.

Nicolas Lopez
University of California Riverside

Date submitted: 30 Oct 2012

Electronic form version 1.4