

Abstract Submitted
for the MAR13 Meeting of
The American Physical Society

Time resolved Magneto-Photoluminescence in $InAs_xP_{1-x}$ alloys¹

TRAVIS MERRITT, MICHAEL MEEKER, GITI A. KHODAPARAST, Virginia Tech, STEPHEN MCGILL, NHMFL, Florida, JOE G. TISCHLER, Naval Research Labs, SUKGEUN G. CHOI, NREL, CHRIS J. PALMSTRØM, Univ. of California Santa Barbara — Recently, g-factor engineering has attracted much attention for potential applications in spintronics. In the case of $InAs_xP_{1-x}$ alloys, a wide range of g-factors, including $g=0$, can be achieved. In order to probe the band-structure and the dynamics of photo-excited carriers in $InAs_xP_{1-x}$ epitaxial films with $x=0.13, 0.4$, we measured NIR absorption spectra at 4K and 300K, as well as magneto-photoluminescence spectra in both the time and frequency domain for magnetic fields in the range of 0-15T and temperatures in the range of 4-90K. From the temporal measurements, we observed strong tunability in the relaxation dynamics as a function of excitation wavelength. We present these experimental observations and compare them with theoretical calculations.

¹Supported by: NSF-Career Award DMR-0846834

Giti Khodaparast
Virginia Tech

Date submitted: 02 Nov 2012

Electronic form version 1.4