

Abstract Submitted
for the MAR13 Meeting of
The American Physical Society

Spin filtering with EuO: Insight from a complex band structure

PAVEL LUKASHEV, Department of Physics and Astronomy & Nebraska Center for Materials and Nanoscience, University of Nebraska, Lincoln, ALEKSANDER WYSOCKI, School of Applied Engineering Physics, Cornell University, JULIAN VELEV, Department of Physics, Institute for Functional Nanomaterials, University of Puerto Rico, MARK VAN SCHILFGAARDE, Department of Physics, Kings College London, England, UK, SITARAM JASWAL, KIRILL BELASHCHENKO, EVGENY TSYMBAL, Department of Physics and Astronomy & Nebraska Center for Materials and Nanoscience, University of Nebraska, Lincoln — Spin-filter tunneling is a promising way to create highly-spin-polarized currents. So far the understanding of the spin-filtering effect has been limited to a free-electron description based on the spin-dependent tunneling barrier height. In this work we employ density-functional calculations to explore the complex bands of EuO as a representative ferromagnetic insulator used in spin-filter tunneling experiments and show that the mechanism of spin filtering deviates significantly from the standard free-electron picture and involves effects associated with the symmetry of spin-dependent evanescent states and the dependence of the decay constant on the transverse wave vector. We demonstrate the importance of the multiorbital band structure with an indirect band gap for spin-filter tunneling. By analyzing the symmetry of the complex bands and the decay rates for different wave vectors and energies we draw conclusions about spin-filter efficiency of EuO. Our results provide a new insight into spin-filter tunneling and may help to design tunnel junctions with enhanced spin polarization.

Pavel Lukashev
Department of Physics and Astronomy & Nebraska Center for
Materials and Nanoscience, University of Nebraska, Lincoln

Date submitted: 05 Nov 2012

Electronic form version 1.4